Formulas for AP Statistics

I. Descriptive Statistics

$$\bar{x} = \frac{\sum x_i}{n}$$

$$s_x = \sqrt{\frac{1}{n-1}\sum(x_i - \bar{x})^2}$$

$$s_p = \sqrt{\frac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{(n_1-1)+(n_2-1)}}$$

$$y = b_0 + b_1 x$$

$$b_1 = \frac{\sum(x_i - x)(y_i - y)}{\sum(x_i - x)^2}$$

$$b_0 = \bar{y} - b_1\bar{x}$$

$$r = \frac{1}{n-1}\sum\left(\frac{x_i - \bar{x}}{s_x}\right)\left(\frac{y_i - \bar{y}}{s_y}\right)$$

$$b_1 = r\frac{s_y}{s_x}$$

$$s_{b_1} = \frac{\sqrt{\dfrac{\sum(y_i - y_i)^2}{n-2}}}{\sqrt{\sum(x_i - \bar{x})^2}}$$

II. Probability

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$E(X) = \mu_X = \sum x_i p_i$$

$$Var(X) = \sigma_X^2 = \sum(x_i - \mu_X)^2 p_i$$

If X has a binomial distribution with parameters n and p, then:

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}$$

$$\mu_X = np$$

$$\sigma_X = \sqrt{np(1-p)}$$

$$\mu_{\hat{p}} = p$$

$$\sigma_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$$

If $\bar{x}$ is the mean of a random sample of size n from an infinite population with mean μ and standard deviation σ, then:

$$\mu_{\bar{x}} = \mu$$

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$$

III.   Inferential Statistics

Standardized test statistic: $\dfrac{\text{estimate - parameter}}{\text{standard deviation of the estimate}}$

Confidence interval: statistic $\pm$ (critical value) $\bullet$ (standard error of statistic)

Single Sample

Statistic	Standard Deviation
Sample Mean	$\dfrac{\sigma}{\sqrt{n}}$
Sample Proportion	$\sqrt{\dfrac{p(1-p)}{n}}$

Two-Sample

Statistic	Standard Deviation
Difference of Sample Means	$\sqrt{\dfrac{\sigma_1^2}{n_1}+\dfrac{\sigma_2^2}{n_2}}$ Special case when $\sigma_1 = \sigma_2$ $\sigma\sqrt{\dfrac{1}{n_1}+\dfrac{1}{n_2}}$
Difference of Sample Proportions	$\sqrt{\dfrac{p_1(1-p_1)}{n_1}+\dfrac{p_2(1-p_2)}{n_2}}$ Special case when $p_1 = p_2$ $\sqrt{p(1-p)}\sqrt{\dfrac{1}{n_1}+\dfrac{1}{n_2}}$

$$\text{Chi-square test statistic} = \sum \dfrac{(\text{observed - expected})^2}{\text{expected}}$$